
AD-A048 529

CLEMSON UNIV S C DEPT OF MATHEMATICAL SCIENCES
A NATURAL APPROACH FOR DETECTING CAUSAL RELATIONSHIP IN TIME SE--ETC(U)
AUG 76 F W MORGAN, M E IRELAND
NO0014-75-C-0451
NB3

LOF |
A048529
A048529
BU

END
PATE
FILMED
2-78
DOC



Frederick W. Morgan, Clemson University M. Edwin Ireland, Clemson University

DDC DECEMBER JAN 5 1978

I. Introduction

A post hoc data analysis in hopes of identifying causal relationships between variables is one that should be regarded with a great deal of suspicion. One need merely consider the etymology of the word "malaria" to realize that unless all factors, or variables, are taken into consideration or experimentally controlled erroneous conclusions concerning cause and effect may be reached.† It is for this reason that statisticians prefer to speak in terms of "predictability" or "correlation" rather than "causation".

In economics, however, a great deal of research has recently been published concerning causal realtionships between economic variables. The basis for this research stems from a definition of causality given by C. W. J. Granger (1964). Basically this definition states that if one can predict the future values of one variable X, using all past values of all other variables in this system better than when predicting future X_t using all past values of all other variables in this system except those of Y, then Y causes X. It is important to note that this definition contains a universe of variables related in some system. In reality, of course, it is impossible to measure all the factors in the system. Hence it may appear that Y_t is causing X_t when indeed a third variable is causing both X_t and Y_t . Granger is careful to point this out in his discussion of the definition.

The inability to measure all variables in the universe certainly exists in economics. However, when an economic theory indicates that a change in one variable should cause a change in another, it may be necessary to identify a statistical relationship between these variables, a relationship which could quite possibly be causal but could not be proven. Thus, though some critics may argue the use of the word "causality", we shall use this term with the understanding of what it connotes in this situation. Wholesale use of techniques for identifying casuality without sound economic reasons for justifying such techniques should be avoided.

Granger's definition of causality is important in that not only does it appeal to our intuition as a reasonable definition, it also lends itself to statistical analyses for causality detection. The appropriate analysis to be used, though, is not apparent. In the next section we propose some analyses that seem natural in this situation but which should not be used because of certain deficiencies. The elimination of these deficiencies leads to what we propose for causality which is given in the fourth section. This discussion will consist of a universe of only two variables, however, the extension to more than two variables follows readily.

II. The Process of Elimination

In Granger's definition one is concerned with being able to predict one variable from past values of itself and another variable. As a criterion of goodness of the prediction Granger employs mean squared error. Thus one's initial attempt at determining if Y causes X would be to regress present X on past X and Y according to

 $x_{t} = a_{1}^{1} x_{t-1} + \dots + a_{n}^{1} x_{t-n} + b_{1}^{1} x_{t-1} + \dots + b_{m}^{1} x_{t-m} + \epsilon_{t}^{1},$ (2.1)

where ε_t -NID(0, σ^2). A significant F-ratio for testing H_: b_i = 0, i = 1, ..., m, would imply that Y_t causes X_t. This result, though, would surely be suspect because in most economic time series a definite trend exists with respect to time. This trend in time present in both series could manifest itself in the significance of the coefficients in the Y_t series when indeed Y_t does not cause X_t.

The next consideration then would be the removal of or an adjustment for the time variable. In the terminology of time series analysis our aim becomes the transformation of the original series to stationary time series. We use the term "stationary" to describe a time series with constant mean having autocovariance a function only of the lag between observations. This condition on the autocovariance function implies also constant variance. Departure from constant variance may be corrected via some transformation as the logarithm. As indicated by Box and Jenkins (1976) stationarity with respect to the mean can be achieved by suitably differencing the individual series involved. For example, if a linear trend is present in the data then one would consider the series $U_t = X_t - X_{t-1}$. Had the trend had a single periodic component then a second difference would be required, namely, U' = U - U - 1.

We return then to the regression model (2.1) where we realize now that the X_t and Y_t are the results of the differencing necessary to produce stationarity. New problems arise, however, that cause the results of a classical regression analysis based on the F-test to be suspect. The standard assumptions for the F-test are that the values of the dependent variable are normally and independently distributed and that the values of the independent variables are fixed. In our situation the values of the "dependent" variable are not necessarily independent (and, in fact, are most often correlated). Furthermore

more than two variables follows readily.

[See 1473]

Reprinted from the 1976 Business and Economic Statistics Section Proceedings of the American Statistical Association Approved for public release; Distribution Unlimited DDC

the lack an ependence is not due to correlated errors (which could be adjusted for) but due to the fact that the "independent" variables are not fixed but random. In fact the "dependent" variable at time t is one of the "independent" variables at some future time t + k. It is this structure that arouses questions concerning the distribution of the regression statistic and thus the validity of the test.

It should also be noted that this regression technique at this point fails for another important reason. It has been shown [Jenkins-Watts, p. 338, (1968)] that even if X_t and Y_t are independent processes having autoregressive structure that spurious cross-correlations are possible. These spurious cross-correlations could manifest themselves in significant values for some of the Y_t coefficients in (2.1) implying causality when in fact none exists.

III. A More General Model

Since the regression model (2.1) has been shown to have serious deficiences in the ensuing analyses, we next consider two different models and the relationship between them. The models to be considered are the univariate autoregressivemoving average representation for each of two series, X, and Y, and the bivariate autoregressive representation of X_t and Y_t jointly. The series X_t and Y_t are assumed to be appropriately transformed or differenced so that each series is covariance stationary. Furthermore, we shall assume that the true causal relationship has remained in the residuals X and Y, although this assumption may be sensitive to the nature of the prefiltering (especially when nonlinear transformations to achieve stationarity are used). The univariate models are:

$$\phi_1(B) \ X_t = \theta_1(B) \ u_t$$

$$\phi_2(B) \ Y_t = \theta_2(B) \ v_t$$
(3.1)

where ϕ_1 , ϕ_2 , θ_1 , and θ_2 are finite degree polynomials in B (where $\mathbf{B}^{\mathbf{K}}\mathbf{X}_{\mathbf{t}}=\mathbf{X}_{\mathbf{t}-\mathbf{k}}$) having constant term I and $\mathbf{u}_{\mathbf{t}}$ and $\mathbf{v}_{\mathbf{t}}$ are white noise processes. The processes are strictly autoregressive if $\theta_1(\mathbf{B})=1$, i=1, 2, and strictly moving averages if $\phi_1(\mathbf{B})=1$, i=1, 2. The assumption that $\mathbf{X}_{\mathbf{t}}$ and $\mathbf{Y}_{\mathbf{t}}$ are stationary implies that the roots of $\phi_1(\mathbf{Z})=0$ lie outside the unit circle. We can subsequently rewrite (3.1) as

$$x_{t} = \phi_{1}^{-1}$$
 (B) θ_{1} (B) $u_{t} = \theta_{1}$ (B) u_{t}
 $y_{t} = \phi_{2}^{-1}$ (B) θ_{2} (B) $v_{t} = \theta_{2}$ (B) v_{t} , (3.2)

where ϕ_1^{-1} (B) and ϕ_2^{-1} (B) are infinite series in B provided they are not identically 1.

The bivariate autoregressive representation is given by

$$\begin{bmatrix} P(B) & Q(B) \\ R(B) & S(B) \end{bmatrix} \begin{bmatrix} X_t \\ Y_t \end{bmatrix} = \begin{bmatrix} a_t \\ b_t \end{bmatrix}$$
 (3.3)

where P, Q, R, and S are polynomials of the form

$$A(B) = \sum_{k=0}^{\infty} a_k B^k$$

with the constant term in P and S strictly 1 and \mathbf{a}_{t} and \mathbf{b}_{t} are white noise series with

$$E\left[\begin{bmatrix} a_t \\ b_t \end{bmatrix} \begin{bmatrix} a_{t+k} & b_{t+k} \end{bmatrix}\right] = \begin{bmatrix} \int (diagonal), k=0 \\ \phi & , k \neq 0 \end{bmatrix}$$

The series are stationary if the roots of

$$\begin{array}{c|c} P(Z) & Q(Z) \\ R(Z) & S(Z) \end{array} = 0$$

lie outside the unit circle.

The relationship between these two representations can be given in terms of their white noise innovations. From (3.2) we can rewrite (3.3) as

$$\begin{bmatrix} P(B) & Q(B) \\ R(B) & S(B) \end{bmatrix} & \begin{bmatrix} \Theta_1(B) & O \\ O & \Theta_2(B) \end{bmatrix} & \begin{bmatrix} u_t \\ v_t \end{bmatrix} = \begin{bmatrix} a_t \\ b_t \end{bmatrix}$$

or

$$\begin{bmatrix} P(B) & \Theta_1(B) & Q(B)\Theta_2(B) \\ R(B) & \Theta_1(B) & S(B)\Theta_2(B) \end{bmatrix} \begin{bmatrix} u_t \\ v_t \end{bmatrix} = \begin{bmatrix} a_t \\ b_t \end{bmatrix}$$

which with substitutions of the form $P^*(B) = P(B) \Theta_1(B)$ yields

$$\begin{bmatrix} P^*(B) & Q^*(B) \\ R^*(B) & S^*(B) \end{bmatrix} \begin{bmatrix} u_t \\ v_t \end{bmatrix} = \begin{bmatrix} a_t \\ b_t \end{bmatrix}$$

Expressing
$$\begin{bmatrix} u_t \\ v_t \end{bmatrix}$$
 in terms of the $\begin{bmatrix} a_t \\ b_t \end{bmatrix}$ yields

$$\begin{bmatrix} u_t \\ v_t \end{bmatrix} = \begin{bmatrix} F(B) & S^*(B) & -F(B) & Q^*(B) \\ -F(B) & R^*(B) & F(B) & P^*(B) \end{bmatrix} \begin{bmatrix} a_t \\ b_t \end{bmatrix}$$

which in single equation becomes

$$u_t = F(B) S^*(B) a_t - F(B) Q^*(B) b_t$$

 $v_t = -F(B) R^*(B) a_t + F(B) P^*(B) b_t$, (3.4)

where $F(B) = 1/\{P^*(B) S^*(B) - Q^*(B) R^*(B)\}$. Thus u_t and v_t are related to one another via a_t and b_t according to (3.4). It is this pair of equations that can be used to determine the causal relationship between u_t and v_t , that is, between X_t and Y_t .

If no causal relationship exists between X_t and Y_t , then Q(B) = R(B) = 0 and (3.4) reduces to

$$u_t = a_t$$

$$v_t = b_t$$

that is, the white noise innovations u_t and v_t have cross correlation zero for all lags. Under a null hypothesis of no causality and assuming that a_t and b_t are Gaussian, a regression of u_t on past, present, and future v_t based on the standard test statistic which does indeed follow an F distribution should find no significant coefficients.

In the case of unidirectional causality, say Y_t causing X_t , R(B) = 0. Substituting these into (3.4) yields

$$u_{t} = \frac{1}{P^{*}(B)} a_{t} - \frac{Q^{*}(B)}{P^{*}(B) S^{*}(B)} b_{t} = A_{t} + B_{t}$$

$$v_{t} = b_{t} . \qquad (3.5)$$

Assuming for now no instantaneous causality, the operator $\frac{Q^*(B)}{p^*(B) S^*(B)}$ is a polynomial in B having constant term zero. Thus u_t is a function of past v_t . Rewriting the first equation in (3.5) to express v_t in terms of u_t produces

$$v_t = -\left[\frac{Q^*(B)}{P^*(B) S^*(B)}\right]^{-1} u_t + \frac{S^*(B)}{Q^*(B)} a_t.$$
 (3.6)

The operator on u_t will contain terms of the form B^{-m} , m>0 (as well as B^{m} terms perhaps), implying that v_t will be a function of future u_t (as well as possibly present and past v_t). A forward stepwise regression finding u_t a function of only past v_t and v_t a function of future and possibly present and past u_t would thus indicate that Y_t causes X_t .

A closer look at (3.5) might arouse some concern about an ordinary least square regression of \mathbf{u}_t on \mathbf{v}_t . In general, the first term \mathbf{A}_t is a moving average in \mathbf{a}_t and the second term \mathbf{B}_t is a moving average in \mathbf{b}_t . Thus each term represents an autocorrelated series. The nature of this autocorrelation is such that when \mathbf{A}_t is correlated ρ_k at lag k the second term \mathbf{B}_t must have autocorrelation $-\rho_k$ at lag k in order that

u be white noise. In a regression u on v it is clear that A represents the error in the regression. It is well known that such autocorrelation among the error terms can result in biased estimates of the variances of the least squares estimator of the regression coefficients (see Appendix A). This causes concern in that one might tend to reject true null hypotheses more often than the significance level of the test would indicate or one might tend to not reject a false null hypothesis when indeed he should. As indicated in Appendix A the nature of the autocorrelation of the "independent" variable determines which of the above is the situation. In the case of unidirectional causality when using a forward stepwise regression, it is easy to see that at each step the next possible "independent" regression variable to enter the model is indeed uncorrelated. In this case the estimator of the error variance is asymptotically unbiased and for small samples the bias should be negligible (our simulations using around 80 observations have indicated no consistent bias in estimates of the error variance). Hence the use of ordinary least squares in a forward stepwise regression is valid.

In the case where Y_t causes X_t (or X_t causes Y_t) and instantaneous causality is present, the polynomials $Q^*(B)$, $P^*(B)$, and $S^*(B)$ each have constant term 1. Thus u_t will be a function of present and past v_t and similarly in (3.6) v_t a function of present and past u_t . Hence the criterion for instantaneous causality. (It should be noted that this criterion only identifies the existence of instantaneous causality. Whether X_t causes Y_t , Y_t causes X_t , or

instantaneous feedback cannot be determined.)
When feedback is present in the system,
both Q(B) and R(B) are nonzero polynomials in B
with constant term zero (assuming no instantaneous causality). Equation (3.4) may now be expressed as

$$u_t = A_1(B) a_t + B_1(B) b_t$$


 $v_t = A_2(B) a_t + B_2(B) b_t$

where A_1 and B_2 have constant term 1 and A_2 and B_1 have constant term 0. Considering u_t as a linear combination of the v_t implies

$$u_t = A_1^{(B)}A_2^{-1}_{(B)}v_t +$$

$$[B_1^{(B)} - A_1^{(B)}A_2^{-1}_{(B)}B_2^{(B)}] b_t$$

Now $A_1(B)A_2^{-1}$ contains negative powers of B as well as positive powers. Thus u_t is a function of past, future, and possible present values of v_t . A symmetric argument holds for v_t as a function of u_t . Hence the criterion for determining feedback in the system. As in the unidirectional



case the absence of autocorrelation in the "independent" regressor variables substantiates the validity of ordinary least squares for the regressions.

IV. The Resulting Procedure

In summary, the procedure that has arisen naturally from the earlier discussion is outlined as follows:

- Individually transform (e.g., taking the natural logarithm of the observations) or difference (e.g., taking the first difference to remove a linear trend) each time series to yield covariance stationary residual series denoted by X_t and Y_t.
- Individually model each series, X_t and Y_t, as an autoregressive-moving average process according to the techniques developed by Box and Jenkins (1976).
- Filter X_t and Y_t according to their estimated models to yield the white noise innovations u_t and v_t respectively.
- 4. Use a forward stepwise regression of u_t on future, present, and past values of v_t and a similar regression of v_t on u_t .
- 5. Determine the causal relationship as follows: a. No causality, -- u_t is not a function of any of the future, present, or past values of v_t nor is v_t a function of any such u_t.
 - b. Y_t causes X_t (and similarly for X_t causing Y_t) but not instantaneously -- u_t is a function only of past v_t and v_t is a function of future and possibly present and past u_t.
 - c. Instantaneous causality -- u_t is a function of only present and past v_t and v_t is a function of only present and past u_t .
 - d. Bi-directional causality -- u_t is a function of future, present, and past v_t and similarly v_t a function of future, present, and past u_t.

We advocate this procedure because (1) it generalizes to the multivariate case and therefore serves as a basis for the detection of causality in systems of two or more variables and (2) the use of a stepwise regression serves not only to determine the direction of causality but also to determine the causality lag. The latter is an important step in system identification. For example, consider that unidirectional causality has been detected from Y_t to X_t . From our regression analysis we have estimated $-\frac{Q^*(B)}{P^*(B) \ S^*(B)} \ \text{in (3.5)} \ \text{and from our original} \ \text{modelling we have estimated } \theta_1(B) \ \text{and } \theta_2(B)$. A Box-Jenkins analysis of the residual series A_t

YOURS YE SEALMEN NOTES THE

will yield an estimate of $P^*(B)$ and hence P(B). Since $S^*(B) = 1$ an estimate for $Q^*(B)$ can be obtained and hence Q(B). From this analysis, then, we can estimate the original model

$$\begin{bmatrix} P(B) & Q(B) \\ 0 & S(B) \end{bmatrix} \begin{bmatrix} X_t \\ Y_t \end{bmatrix} = \begin{bmatrix} a_t \\ b_t \end{bmatrix}$$

(knowing that R(B) = 0 in this case).

V. Applications of the Procedure

In order to demonstrate the application of this procedure in bivariate situations the results of a simulation and an analysis of two economic times series are given. The results concerning the economic time series are quite interesting in that they differ from any other results yet published.

For the simulation 88 observations were generated for each of two series $X_{\mbox{\scriptsize t}}$ and $Y_{\mbox{\scriptsize t}}$ according to

$$X_{t} = .7X_{t-1} + Y_{t-2} + a_{t}$$
, $a_{t}^{-NID(0,1)}$
 $Y_{t} = .75Y_{t-1} - .5Y_{t-2} + b_{t}$, $b_{t}^{-NID(0,1)}$.

Due to the choice of coefficients the processes are stationary. A Box-Jenkins modelling of the series individually yielded

$$x_t = .975x_{t-1} - .297x_{t-2} + u_t - .257_{t-8}$$

 $y_t = .850y_{t-1} - .465y_{t-2} + v_t$

Upon filtering each series to yield u_t and v_t and performing forward stepwise regressions of u_t on v_t and v_t on u_t , the models obtained were

$$u_t = .85v_{t-2} + .28v_{t-3} + .27v_{t-4}$$
 $- .35v_{t-5} + error$
 $(F_{4,64} = 11.25 R^2 = .41 D.W. = 2.61)$
 $v_t = .36u_{t+2} + .16u_{t+3} + error$
 $(F_{2,66} = 21.03 R^2 = .39 D.W. = 2.47)$

These results indicate causality from Y_t to X_t . It is also interesting to note the magnitude of the Durbin-Watson statistics indicating the anticipated serial correlation for the error terms.

Applying our technique to quarterly observations of the money supply and GNP for the period 1953 through 1974, we arrived at the following model for GNP:

$$G_{t} = (1-B) \ln (GNP)$$

 $(G_{t} - .007) = .449(G_{t-1} - .007) + E_{t}$
 $G_{t} = E_{t} = (G_{t} - .007) - .449(G_{t-1} - .007)$.

That is the first differences of the natural log of GNP, which is approximately equivalent to the rate of growth from quarter to quarter (a transformation necessary to achieve stationarity) can be modeled as a simple first order autoregressive (AR) process.

For money (M1), the model is

$$M_{t} = (1 - B) \ln (M1)_{t}$$

$$(M_{t} - .004) = .63(M_{t-1} - .004) + Z_{t} - .259Z_{t-10}$$

$$m_{t} = Z_{t} = (M_{t} - .004) + .63(M_{t-1} - .004) + .259Z_{t-10}$$

That is, the first differences of the natural logs of money is described by an autoregressive – moving average (ARMA) model. The residuals of these correctly filtered series, g_{t} and m_{t} , can now be used in a regression to test for

Using a forward selection regression procedure combined with the properly prefiltered values of money and GNP, we arrived at the following models:

$$g_t = .49_{mt-2} + .68m_{t-3} + .42m_{t+1} - .54m_{t+3}$$

$$(.1850) \quad (.1874) \quad (.1831) \quad (.1880)$$
 $R^2 = .2775 \quad F(4,65) = 6.2408 \quad D.W. = 2.41$

and

$$m_t = -.1359g_{t-3} + .1901g_{t+3}$$
(.0649) (.0647)
 $R^2 = .1573 \quad F(2,67) = 6.2520 \quad D.W. = 2.1097$,

where the values in parentheses below the coefficients are the standard errors of the regression coefficients. In both regressions, the $\rm R^2$'s are small but still significant at the .05 level. Of course, when evaluating this $\rm R^2$, it must be remembered that the values used in the regression were the residuals of previous models. Hence, the explanatory power of these models is actually quite good. Since coefficients of future values appear in both regressions significantly different from zero, we have identified a bidirectional or feedback relationship between money and income.

APPENDIX A

Consider the model

$$y_t = \beta x_t + z_t$$

where x_t and z_t are zero mean processes with z_t

having variance σ_z^2 and autocorrelation ρ_k . The variance of the ordinary least squares estimator b of β given by

$$\mathsf{b} = (\underline{\mathsf{x}}'\underline{\mathsf{x}})^{-1}\underline{\mathsf{x}}'\underline{\mathsf{y}}$$

is

$$Var (b) = (x'x)^{-1} x' Vx(x'x)^{-1}$$

where

$$\mathbf{v} = \sigma_{\mathbf{z}}^{2} \begin{bmatrix} 1 & \rho_{1} & \rho_{\mathbf{z}} & \cdots & \rho_{n-1} \\ \rho_{1} & 1 & \rho_{1} & \cdots & \rho_{n-2} \\ \vdots & \vdots & \vdots & \vdots \\ \rho_{n-1} & \rho_{n-2} & \rho_{n-3} & \cdots & 1 \end{bmatrix}$$

It is easy to show that

$$Var (b) = \frac{\sigma_z^2}{\sum_{i=1}^{n} x_i^2} \left\{ 1 + 2\rho_1 \frac{\sum_{i=1}^{n-1} x_i x_i + 1}{\sum_{i=1}^{n} x_i^2} + 2\rho_2 \frac{\sum_{i=1}^{n-2} x_i x_i + 2}{\sum_{i=1}^{n} x_i^2} \right\}$$

$$2\rho_{n-1}\frac{x_1x_n}{\sum_{i=1}^n x_i^2}$$

$$= \frac{\sigma_2^2}{\sum_{i=1}^n x_i^2} K$$

If $\sigma_{\bf z}^{-2}$ were known, then the usual least squares estimator of the variance $\frac{\sigma_{\bf z}^{-2}}{\sum\limits_{i=1}^n x_i^{-2}}$. Hence if the

 $\mathbf{x_t}$ are autocorrelated with the same sign as the ρ_k then the ordinary least squares estimator will underestimate the true variance of b; if autocorrelated with opposite sign, the estimator will overestimate the variance of b.

Similarly it can be shown that the residual sum of square for least squares estimator has expectation (n-K) σ_s^2 .

In estimating $\sigma_{\mathbf{z}}^{-2}$ the same relationships hold as above accentuating the bias in estimating the variance of b using ordinary least squares.

It should be noted, however, that if the x_t are uncorrelated, the affect of the autocorrelated errors will likely be neglible. In fact in this case the estimators are asymptotically unbiased.

the only known carrier is the anopheles mosquito.

REFERENCES

- Box, G. E. P. and Jenkins, G. M., Time Series
 Analysis: Forecasting and Control, Revised
 edition. San Francisco: Holden-Day, 1976.
- Granger, C. W. J., "Processes Involving Feedback"
 (Chapter 7) in <u>Spectral Analysis of Economic</u>
 <u>Time Series</u> (with M. Hatanaka). Princeton,
 New Jersey: Princeton University Press, 1964.
- Jenkins, G. M. and Watts, D. G., Spectral Analysis and its Applications. San Francisco: Holden-Day, 1968.

[†]The word malaria means "bad air". This term came into usage because it was thought that people contracted the disease from the air present in swampy, lowlying areas when in fact

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered) READ INSTRUCTIONS REPORT DOCUMENTATION PAGE BEFORE COMPLETING FORM 2. GOVT ACCESSION NO. 3. RECIPIENT'S CATALOG NUMBER REPORT NUMBER N83 S. TYPE OF REPORT & PERIOD COVERED TITLE (and Subtitle) A Natural Approach for Detecting Causal Technical Report, Relationships in Time Series, PERFORMING ORG. REPORT NUMBER . AUTHOR(a) S. CONTRACT OR GRANT NUMBER(s) Nggg14-75-C-9451 Frederick W. / Morgan M. Edwin / Ireland PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PERFORMING ORGANIZATION NAME AND ADDRESS Clemson University Dept. of Mathematical Sciences NR 042-271 Clemson, South Carolina 29631 11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research 23 Code 436 Arlington, Va. 14. MONITORING AGENCY NAME & ADDRESS(II different from Controlling Office) 18. SECURITY CLASS. Unclassified 15a. DECLASSIFICATION/DOWNGRADING 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) 18. SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary and identity by block number) Stepwise Regression Causality Autocorrelation Time Series Stationarity Bivariate Autoregressive Process 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A definition of causality given by C.W.J. Granger lends itself to statistical hypothesis testing for identifying the causal relationship between two time series. The appropriate statistical analysis to be used, however, is not apparent from the definition. One's first impression is to regress present values of one time series on past values of both time series. There are, however, several weaknesses inherent in this procedure Natural modifications of this procedure lead eventually to a Box-Jenkins approach for suitably preprocessing the time series followed by a regression analysis of the residuals, an analysis which identifies the causal structure between the two the residuals, an analysis which identifies the causal time series. DD 1 JAN 73 1473 EDITION OF I NOV 65 IS OBSOLETE

407183

UNCLASSIFIED
SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)